


Composition Analysis of Fully Aromatic Polyester by Py-GC Utilizing Reactive Pyrolysis

In analysis of polyesters and other condensation polymers by the Py-GC technique, reactive pyrolysis in the presence of tetramethyl ammonium hydroxide (TMAH) is very useful. Ester bonds of fully aromatic polyesters, in particular, are almost completely hydrolyzed with this method to give corresponding methylesters quantitatively. For example, Fig. 1 shows a pyrogram of reactive pyrolysis of a fully aromatic polyester made from p-hydroxy benzoic acid (PHB), terephthalic acid (TA) and 4,4-bisphenol (BP). Only the peaks of methyl derivatives of monomers are observed, indicating that the polyester is completely hydrolyzed into monomers. Fig. 2 shows the analysis result of another polyester prepared from the same monomers at different feed ratios. It is evident that the result is in a good agreement with the feed ratios. These results demonstrate that Py-GC utilizing reactive pyrolysis is a particularly useful technique for composition analysis of fully aromatic polyesters.



Keyword : Aromatic Polyester, Composition Analysis, Reactive Pyrolysis, TMAH

Application: General Polymer Analysis

Please forward your inquiries via our web page at: (http://www.frontier-lab.com/), or send us a fax message.

R&D and manufactured by:

Frontier Laboratories Ltd.

1-8-14, Saikon, Koriyama, Fukushima, 963-8862 Japan

Phone: 81-24-935-5100 Fax: 81-24-935-5102

® : Registered trademark of Frontier Laboratories Ltd.

Your dealer: